

**LIST OF BAN-LIFTED PLANT ITEMS AND KEY CONDITIONS (MEET THE STANDARDS SET BY THE MINISTER OF AGRICULTURE, FORESTRY AND FISHERIES)**

1. April, 2010

COUNTRIES / DISTRICTS	FRESH FRUITS		BAN-LIFTED YEAR	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE																														
	ITEMS	CULTIVARS																																		
UNITED STATES OF AMERICA	HAWAIIAN ISLANDS	PAPAYA	SOLO	April, 1969	①MEDITERRANEAN FRUIT FLY ②ORIENTAL FRUIT FLY COMPLEX ③MELON FLY	Vapor Heat Treatment: The fruits shall be disinfested by using saturated vapor at vapor heat treatment facilities till the temperature of the innermost fruit pulp reaches 47.2°C.	SHIP CARGO AIR CARGO AIR HAND BAGGAGE																													
		MANGO	KEITT HADEN	May, 2000																																
	EXCLUDING HAWAIIAN IS.	CHERRY	1.BING 2.LAMBERT 3.VAN 4.RAINIER 5.GARNET 6.TULARE 7.BROOKS 8.LAPIN 9.SWEET HEART 10.CHELAN 11.EXCEPT ABOVE	1.BING 2.LAMBERT 3.VAN 4.RAINIER 5.GARNET 6.TULARE 7.BROOKS 8.LAPIN 9.SWEET HEART 10.CHELAN 11.EXCEPT ABOVE	①January, 1978	CODLING MOTH	Methyl Bromide Fumigation ① Of cultivars 1~10 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>32g/m<sup>3</sup></td> <td>22°C or above</td> <td>2 hours</td> <td>50% or less</td> </tr> <tr> <td>40g/m<sup>3</sup></td> <td>17~22°C or below</td> <td>2 hours</td> <td>50% or less</td> </tr> <tr> <td>48g/m<sup>3</sup></td> <td>12~17°C or below</td> <td>2 hours</td> <td>50% or less</td> </tr> <tr> <td>64g/m<sup>3</sup></td> <td>6~12°C or below</td> <td>2 hours</td> <td>50% or less</td> </tr> </tbody> </table> ② Or all the cultivars including cultivars 1~10, and 11 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration exposure</th> </tr> </thead> <tbody> <tr> <td>64g/m<sup>3</sup></td> <td>6~12°C or below</td> <td>2 hours or more</td> <td>61.9 or up</td> </tr> </tbody> </table> 1. Plants and areas: The cherry fresh fruits shall be produced in designated areas by American plant protection authorities to be subject to intensive trapping surveys and fresh fruit surveys for Codling moths as the result of the surveys set forth in Article 2. 2. Survey in designated production areas Trapping survey (traps shall be set at the density of one per 1 hectare). ① Trapping survey ② Fresh fruit survey 3. Inspection and Certification in Producing Area The phytosanitary certificate shall include the following statements: (a) The fruit is not infested with Codling moths. (b) The fruit has been produced at the designated area where the number of Codling moths in the survey do not exceed 10 per trap per week on average in California and 30 per trap per week on average in Oregon and Washington.	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	32g/m <sup>3</sup>	22°C or above	2 hours	50% or less	40g/m <sup>3</sup>	17~22°C or below	2 hours	50% or less	48g/m <sup>3</sup>	12~17°C or below	2 hours	50% or less	64g/m <sup>3</sup>	6~12°C or below	2 hours	50% or less	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure	64g/m <sup>3</sup>	6~12°C or below	2 hours or more	61.9 or up	SHIP CARGO AIR CARGO
					Dosage			Temperature	Duration of exposure	Ratio of fruit capacity																										
					32g/m <sup>3</sup>			22°C or above	2 hours	50% or less																										
					40g/m <sup>3</sup>			17~22°C or below	2 hours	50% or less																										
					48g/m <sup>3</sup>			12~17°C or below	2 hours	50% or less																										
					64g/m <sup>3</sup>			6~12°C or below	2 hours	50% or less																										
					Dosage			Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure																										
					64g/m <sup>3</sup>			6~12°C or below	2 hours or more	61.9 or up																										
②December, 1986																																				
May, 1992																																				
January, 1995																																				
October, 1996																																				
July, 1999																																				
May, 2001																																				
October, 2001																																				
Jun, 2009																																				
ALL VARIETY																																				
ALL VARIETY																																				
EXCLUDING HAWAIIAN IS.	WALNUT IN SHELL	1.HARTLEY 2.PAYNE 3.FRANQUETTE	April, 1986		Methyl bromide fumigation: <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of stone capacity</th> </tr> </thead> <tbody> <tr> <td>56g/m<sup>3</sup></td> <td>15.6°C or above</td> <td>4 hours</td> <td>55% or less</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of stone capacity	56g/m <sup>3</sup>	15.6°C or above	4 hours	55% or less																							
Dosage	Temperature	Duration of exposure	Ratio of stone capacity																																	
56g/m <sup>3</sup>	15.6°C or above	4 hours	55% or less																																	
EXCLUDING HAWAIIAN IS.	NECTARINE	1.SUMMER GRAND 2.SPRING RED 3.FIR EBRITE 4.FANTASIA 5.MAY GRAND 6.RED DIAMOND 7.MAY FINE 8.MAY GLO 9.MAY DIAMOND 10.ROYAL GIANT 11.EXCEPT ABOVE	1.SUMMER GRAND 2.SPRING RED 3.FIR EBRITE 4.FANTASIA 5.MAY GRAND 6.RED DIAMOND 7.MAY FINE 8.MAY GLO 9.MAY DIAMOND 10.ROYAL GIANT 11.EXCEPT ABOVE	June, 1988		Methyl bromide fumigation: <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>48g/m<sup>3</sup></td> <td>21°C or above</td> <td>2 hours</td> <td>50% or less</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	48g/m <sup>3</sup>	21°C or above	2 hours	50% or less																						
				Dosage			Temperature	Duration of exposure	Ratio of fruit capacity																											
				48g/m <sup>3</sup>			21°C or above	2 hours	50% or less																											
				May, 1993																																
				May, 1995																																
				August, 2000																																
August, 2000																																				
August, 2000																																				
EXCLUDING HAWAIIAN IS.	EUROPEAN PLUM	1.D'AGEN 2.TULARE GIANT 3.MOYER 4.EXCEPT ABOVE	①April, 2001 ②August, 2005		Methyl bromide fumigation: ① Of cultivars 1~3 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>48g/m<sup>3</sup></td> <td>20°C or above</td> <td>2 hours</td> <td>50% or less</td> </tr> </tbody> </table> ② Or of all the cultivars including cultivars 1~3 and 4 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration exposure</th> </tr> </thead> <tbody> <tr> <td>48g/m<sup>3</sup></td> <td>20°C or above</td> <td>2 hours or more</td> <td>72.1 or up</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	48g/m <sup>3</sup>	20°C or above	2 hours	50% or less	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure	48g/m <sup>3</sup>	20°C or above	2 hours or more	72.1 or up	SHIP CARGO AIR CARGO														
Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																																	
48g/m <sup>3</sup>	20°C or above	2 hours	50% or less																																	
Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure																																	
48g/m <sup>3</sup>	20°C or above	2 hours or more	72.1 or up																																	
①WASHINGTON, OREGON AND CALIFORNIA  ②EXCLUDING HAWAIIAN IS.	APPLE	1.RED DELICIOUS 2.GOLDEN DELICIOUS 3.FUJI 4.BRAEBURN 5.GRANNY SMITH 6.GALA 7.JONA GOLD 8.EXCEPT ABOVE	1.RED DELICIOUS 2.GOLDEN DELICIOUS 3.FUJI 4.BRAEBURN 5.GRANNY SMITH 6.GALA 7.JONA GOLD 8.EXCEPT ABOVE	August, 1994	①CODLING MOTH ②FIRE BLIGHT	After the following A treatment, the B treatment shall be done: A Cold Treatment: After the fruit pulp temperature reaches 2.2 °C, the fruits shall be disinfested by the same temperature (2.2°C) for 55 consecutive days at cold treatment facilities. B Methyl Bromide Fumigation: ① Of cultivars 1~7 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>56g/m<sup>3</sup></td> <td>10°C or above</td> <td>2 hours</td> <td>50.9% or less</td> </tr> </tbody> </table> ② Or of all the cultivars including cultivars 1~7 and 8 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration exposure</th> </tr> </thead> <tbody> <tr> <td>56g/m<sup>3</sup></td> <td>10°C or above</td> <td>2 hours or more</td> <td>85.5 or up</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	56g/m <sup>3</sup>	10°C or above	2 hours	50.9% or less	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure	56g/m <sup>3</sup>	10°C or above	2 hours or more	85.5 or up	SHIP CARGO AIR CARGO													
				Dosage			Temperature	Duration of exposure	Ratio of fruit capacity																											
				56g/m <sup>3</sup>			10°C or above	2 hours	50.9% or less																											
				Dosage			Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration exposure																											
56g/m <sup>3</sup>	10°C or above	2 hours or more	85.5 or up																																	
July, 1999																																				
October, 2001																																				
October, 2001																																				
EXCLUDING HAWAIIAN IS.	TOMATO		April, 1997 September, 1999	TABACCO BLUE MOLD																																
ARGENTINE REPUBLIC	GRAPE FRUIT		April, 2003	MEDITERRANEAN FRUIT FLY	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers as follows: (Grape fruit) After the fruit pulp temperature reaches 1.9°C, the fruits shall be disinfested by the temperature below 2.3°C for 19days. Or after the fruit pulp temperature reaches 3.0°C, the fruits shall be disinfested by the temperature below 3.2°C for 23 days. (Sweet orange) After the fruit pulp temperature reaches 1.9 °C, the fruits shall be disinfested by the temperature below 2.2°C for 21days. (Lemon) After the fruit pulp temperature reaches 1.9 °C, the fruits shall be disinfested by the temperature below 2.2°C for 19days. Or after the fruit pulp temperature reaches 3.0°C, the fruits shall be disinfested by the temperature below 3.2°C for 24 days.	SHIP CARGO AIR CARGO																														
	SWEET ORANGE	VALENCIA																																		
	LEMON																																			

**LIST OF BAN-LIFTED PLANT ITEMS AND KEY CONDITIONS (MEET THE STANDARDS SET BY THE MINISTER OF AGRICULTURE, FORESTRY AND FISHERIES)**

1. April, 2010

COUNTRIES / DISTRICTS		FRESH FRUITS		BAN-LIFTED YEAR	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE																																					
		ITEMS	CULTIVARS																																									
STATE OF ISRAEL		SWEET ORANGE	1.SHAMOUTI 2.VALENCIA	June, 1972	MEDITERRANEAN FRUIT FLY	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers as follows: (Orange) After the fruit pulp temperature reaches 0.5°C, the fruits shall be disinfested by the temperature for 14 days. Or after the fruit pulp temperature reaches 1.5°C, the fruits shall be disinfested by the temperature for 16 days. (Grape fruit) After the fruit pulp temperature reaches 0.5°C, the fruits shall be disinfested by the temperature for 13 days. Or after the fruit pulp temperature reaches 1.5°C, the fruits shall be disinfested by the temperature for 16 days. (Sweety) After the fruit pulp temperature reaches 1.5°C, the fruits shall be disinfested by the temperature for 16 days. (Pomelo) After the fruit pulp temperature reaches 1.0°C, the fruits shall be disinfested at 1.5°C or below for 15 days. (Lemon) After the fruit pulp temperature reaches 1.5°C, the fruits shall be disinfested by under 1.5°C for 16 days.  Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches 0.0°C, it has been kept at 0.0°C or below for 12 consecutive days. Or after it reaches 1.1°C, it has been kept at 1.1°C or below for 14 consecutive days.	SHIP CARGO AIR CARGO																																					
		GRAPE FRUIT																																										
		SWEETIE		March, 1990																																								
		POMELO		December, 1998																																								
		LEMON		May, 2008																																								
		PERSIMMON	TRIUMPH	November, 2003																																								
REPUBLIC OF ITALY		SWEET ORANGE	TAROCCO	March, 2005	MEDITERRANEAN FRUIT FLY	The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches 1.2°C, and has been kept at 1.2°C or below for 14 consecutive days.	SHIP CARGO AIR CARGO																																					
INDIA		MANGO	1.ALPHONSO 2.KESAR 3.CHAUSA 4.BANGAN PALLI 5.MALLIKA 6.LANGRA	June, 2006	①ORIENTAL FRUIT FLY COMPLEX ②MELON FLY	Vapor Heat Treatment: In Vapor Heat Treatment facilities, the fruits shall be disinfested in saturated water vapor. The air temperature inside the treatment chamber shall be raised step by step to 50°C or above, so that the pulp temperatures of the center of the fruits shall be raised to 47.5°C and be kept continuously at 47.5°C or above for 20 minutes.	SHIP CARGO AIR CARGO																																					
AUSTRALIA	EXCLUDING TASMANIA	GENUS CITRUS		January, 2005	①MEDITERRANEAN FRUIT FLY ②QUEENSLAND FRUIT FLY	1. Plants and areas Fresh citrus fruits produced in areas of Australia designated by the plant protection authorities of Australia to be subject to intensive trapping surveys and fresh fruit surveys for the Mediterranean fruit flies and Queensland fruit flies set forth in Article 2. 2. Surveys in production areas ① Trapping survey ② Fresh fruit survey 3. Inspection and certification in production areas (a part of extract) The phytosanitary certificate shall include the following statements: (a) The fruit is not infested with the Mediterranean fruit fly or Queensland fruit fly. (b) The plants have been produced in designated areas where no Mediterranean fruit flies and Queensland fruit flies have been detected in the surveys as the result of surveys set forth in Article 2.  Cold Treatment: At cold treatment facilities and refrigerated shipping containers, the fruits shall be cooled until the fruit pulp temperature has reached 1.0°C and maintains it for 16 consecutive days, 2.1°C for 18 consecutive days, and 3.1°C for 20 consecutive days.  Cold Treatment: The fruits shall be disinfested at cold treatment facilities and refrigerated shipping containers at the fruit pulp temperature will maintain 1.0°C for 14 consecutive days, 2.1°C for 16 consecutive days, or 3.1°C for 18 consecutive days.	SHIP CARGO AIR CARGO																																					
		SWEET ORANGE	1.VALENCIA 2.WASHINGTON NAVEL	June, 1982 October, 2001																																								
		IMPERIAL ELLENDALE MURCOTT MINNEOLA		April, 1999 October, 2001																																								
		LEMON		May, 1992 October, 2001																																								
		MANGO	1.KENSINGTON 2.R2E2 3.KEITT 4.KENT 5.PALMAR	October, 1994 December, 1999																																								
		TASMANIA	APPLE	FUJI	December, 1998	CODLING MOTH	Methyl Bromide fumigation: ① Fuji, Jonagold <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>48g/m<sup>3</sup></td> <td>17°C or above</td> <td>2 hours</td> <td>53% or less</td> </tr> </tbody> </table> ② All the cultivars including ① <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration</th> </tr> </thead> <tbody> <tr> <td>48g/m<sup>3</sup></td> <td>17°C or above</td> <td>2 hours or more</td> <td>76.4 or up</td> </tr> </tbody> </table> Methyl bromide fumigation: ① Lapin <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Room temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>50g/m<sup>3</sup></td> <td>12°C or above</td> <td>17°C or above</td> <td>2 hours</td> <td>44.5% or less</td> </tr> </tbody> </table> ② Lapin and other cultivars <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Room temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration</th> </tr> </thead> <tbody> <tr> <td>50g/m<sup>3</sup></td> <td>12°C or above</td> <td>17°C or above</td> <td>2 hours or more</td> <td>77.9 or up</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	48g/m <sup>3</sup>	17°C or above	2 hours	53% or less	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration	48g/m <sup>3</sup>	17°C or above	2 hours or more	76.4 or up	Dosage	Temperature	Room temperature	Duration of exposure	Ratio of fruit capacity	50g/m <sup>3</sup>	12°C or above	17°C or above	2 hours	44.5% or less	Dosage	Temperature	Room temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration	50g/m <sup>3</sup>	12°C or above	17°C or above	2 hours or more	77.9 or up	SHIP CARGO AIR CARGO AIR CARGO AIR HAND BAGGAGE
	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																																								
	48g/m <sup>3</sup>	17°C or above	2 hours	53% or less																																								
	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration																																								
	48g/m <sup>3</sup>	17°C or above	2 hours or more	76.4 or up																																								
Dosage	Temperature	Room temperature	Duration of exposure	Ratio of fruit capacity																																								
50g/m <sup>3</sup>	12°C or above	17°C or above	2 hours	44.5% or less																																								
Dosage	Temperature	Room temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration																																								
50g/m <sup>3</sup>	12°C or above	17°C or above	2 hours or more	77.9 or up																																								
		CHERRY	LAPIN AND OTHER CULTIVARS	March, 2005			SHIP CARGO AIR CARGO																																					
			ALL VARIETY	December, 2008	1. Plants and areas: The cherry fresh fruits shall be produced in designated areas by Australian plant protection authorities to be subject to intensive trapping surveys and fresh fruit surveys for Codling moths as the result of the surveys set forth in Article 2. 2. Survey in designated production areas ① Trapping survey ② Fresh fruit survey 3. Inspection and Certification in Producing Area (a part of excerpts) The phytosanitary certificate shall include the following statements: (a) The fruit shall not be infested with Codling moths. (b) The fruit shall be produced in the designated area where Codling moths have not been trapped, and be produced in the designated producing area where the number of Codling moths in the survey do not exceed seven per trap per week on average as the result of surveys set forth in Article 2. Moreover, it shall be produced in the designated area where no Codling moths have been detected.																																							

**LIST OF BAN-LIFTED PLANT ITEMS AND KEY CONDITIONS (MEET THE STANDARDS SET BY THE MINISTER OF AGRICULTURE, FORESTRY AND FISHERIES)**

1. April, 2010

COUNTRIES / DISTRICTS	FRESH FRUITS		BAN-LIFTED YEAR	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE												
	ITEMS	CULTIVARS																
KINGDOM OF THE NETHERLANDS	TOMATO AND BELL PEPPER		February, 1993	MEDITERRANEAN FRUIT FLY	1. Plants and areas Fruits shall be produced in sites (Designated Production Sites) designated by the Plant Quarantine Authority of The Netherlands as areas where no Mediterranean fruit fly has occurred and satisfy the following conditions: ① The fruits shall be produced at the facilities (Designated Production Facilities) designated by the Plant Quarantine Authority of The Netherlands. ② The fruits shall be produced at the areas (Quarantine Monitoring Areas) designated by the Plant Quarantine Authority of The Netherlands as areas which should be watched the entry of Mediterranean fruit fly as the result of occurrence survey mentioned in Article 2 during the time when Mediterranean fruit flies have been confirmed to be no fear of spreading. 2. The Occurrence Survey in quarantine monitoring areas, designated production sites and designated production facilities. ① Pheromone trap monitoring ② Orchard fruit monitoring 3. Inspection and Certification in Producing Area (a part of excerpts) The phytosanitary certificate shall be mentioned the following additional remarks as the result of the occurrence survey mentioned in Article 2: The fruit shall be produced at the designated production facilities where no Mediterranean fruit flies have not been trapped in the designated producing areas where no Mediterranean fruit flies have not been trapped.	SHIP CARGO AIR CARGO												
	STRAWBERRY, CUCUMBER, EGGPLANT, GRAPE, AUTUMN PUMPKIN AND MELON		February, 1998															
CANADA	CHERRY	LAMBERT	June, 1982	CODLING MOTH	Methyl Bromide Fumigation <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>32g/m<sup>3</sup></td> <td>22°C or more</td> <td>2 hours</td> <td>49% or less</td> </tr> <tr> <td>48g/m<sup>3</sup></td> <td>17~22°C or below</td> <td>2 hours</td> <td>49% or less</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	32g/m <sup>3</sup>	22°C or more	2 hours	49% or less	48g/m <sup>3</sup>	17~22°C or below	2 hours	49% or less	SHIP CARGO AIR CARGO
	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity														
32g/m <sup>3</sup>	22°C or more	2 hours	49% or less															
48g/m <sup>3</sup>	17~22°C or below	2 hours	49% or less															
TOMATO		September, 1996 September, 1999	TABACCO BLUE MOLD															
REPUBLIC OF COLOMBIA	YELLOW PITAYA	(Reference) CACTUS FRUIT	April, 1999	MEDITERRANEAN FRUIT FLY	Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities by using saturated vapor at the fruit pulp temperature of 46.0°C and more for 20 minutes.	SHIP CARGO AIR CARGO												
	MANGO	TOMMY ATKINS	October, 2009															
SPAIN	①LEMON		December, 1988	MEDITERRANEAN FRUIT FLY	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers in the following means: ① Lemon and Clementine After the fruit pulp temperature reaches 2.0°C, the fruits shall be disinfested at the fruit pulp temperature of 2.0°C or below for 16 consecutive days. ② Sweet Orange After the fruit pulp temperature reaches 1.5°C, the fruit shall be disinfested at the fruit pulp temperature of 2.0°C or below for 17 consecutive days.	SHIP CARGO AIR CARGO												
	②CLEMANTINE		January, 2004															
	③SWEET ORANGE	1.NAVEL 2.VALENCIA 3.SALUSTIANA	September, 1996 January, 2004															
KINGDOM OF SWAZILAND	SWEET ORANGE	1.WASHINGTON NAVEL 2.VALENCIA 3.TOMANGO 4.PROTEA	June, 1973	MEDITERRANEAN FRUIT FLY	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches minus 0.6°C and keeps the same temperature for 12 consecutive days.	SHIP CARGO AIR CARGO												
	GRAPE FRUIT CLEMANTINE		June, 2007	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches 0.6°C and keeps the same temperature for 14 consecutive days.														
KINGDOM OF THAILAND	①MANGO	1.NAN-KLARNG-WUN	March, 1987	①ORIENTAL FRUIT FLY COMPLEX ②MELON FLY	Vapor Heat Treatment: ① Mango •Nan-Klarn-g-wun The fruits shall be disinfested by vapor heat treatment facilities, through the use of saturated vapor for 10 minutes at the temperature of 46.5°C or higher at the innermost fruit pulp, or through the use of saturated vapor for 20 minutes at the temperature of 47.0°C at the innermost fruit pulp after a steady increase in innermost temperature to 43.0°C without using vapor . •Nan Dorkmai Pimsen-Daeng,Rad,and Mahachanok The fruits shall be disinfested by vapor heat treatment facilities, through the use of saturated vapor for 20 minutes at the temperature of 47.0°C at the innermost temperature of 43.0°C without using vapor. ② Mangosteen The fruits shall be disinfested by vapor heat treatment facilities through the use of saturated vapor for 58 minutes at a temperature of 46°C or higher at the innermost fruit pulp.	SHIP CARGO AIR CARGO AIR HAND BAGGAGE												
		2.NAM-DORKMAI 3.PIMSEN-DAENG 4.RAD 5.MAHACHANOK	February, 1993 November, 2006															
		②MANGOSTEEN	April, 2003				ORIENTAL FRUIT FLY COMPLEX											
TAIWAN	SWEET ORANGE	1.TANKAN	December, 1975	ORIENTAL FRUIT FLY COMPLEX	(EDB: A means of an alternative treatment is being developed.)	SHIP CARGO AIR CARGO AIR HAND BAGGAGE												
		2.LIUCHENG	April, 1980															
	MANGO	KEITT	June, 1976	①ORIENTAL FRUIT FLY COMPLEX														
	PAPAYA	SOLO	March, 1991	②MELON FLY														
		TAINO No2	December, 2004															
	MANGO	1.IRWIN	June, 1976 March, 1989	ORIENTAL FRUIT FLY COMPLEX	Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities through the use of saturated vapor for 30 minutes at a temperature of 46.5°C or higher at the innermost fruit pulp. And after that, the temperature shall become a normal temperature rapidly.													
		2.HARDEN	March, 1991															
	SWEET ORANGE	PONKAN	November, 1969 March, 1988	Cold Treatment: At cold treatment facilities, after fresh fruit pulp temperature reaches 1°C and keeps the same temperature for 14 consecutive days.														
POMELO		December, 1999	Cold Treatment: At cold treatment facilities, after fresh fruit pulp temperature reaches 1.0°C and keeps the same temperature for 12 consecutive days.															
LITCHI		April, 1980 March, 1988	ORIENTAL FRUIT FLY COMPLEX	After the following A treatment is done, the B treatment will be started. A Vapor Heat Treatment: At vapor heat treatment facilities, to be confirmed that the fruit pulp temperature has increased steadily from 30.0°C to 41.0°C within 45 minutes and maintain the fruit temperature of 46.2°C for 20 minutes. B Cold Treatment: At cold treatment facilities, the fruit pulp temperature is lowered to 2°C, and is being kept for 42 hours. In this case, after the vapor heat treatment of the above A, the fruit pulp temperature is lowered to 2°C within 6 hours.														
	GRAPE	1.KYOHO 2.ITALY		December, 1997	Cold Treatment: At cold treatment facilities, after the fruit pulp temperature reaches 0.5°C and keeps the same treatment at and under 1.0°C for 12 consecutive days.													

**LIST OF BAN-LIFTED PLANT ITEMS AND KEY CONDITIONS (MEET THE STANDARDS SET BY THE MINISTER OF AGRICULTURE, FORESTRY AND FISHERIES)**

1, April, 2010

COUNTRIES / DISTRICTS	FRESH FRUITS		BAN-LIFTED YEAR	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE																
	ITEMS	CULTIVARS																				
PEOPLES REPUBLIC OF CHINA	新疆ウイグル自治区 Xinjiang Uygur Autonomous Region	MELON	(Reference) HAMI MELON 哈密瓜 =HAMI URI	March, 1988	MELON FLY	1. Plants and areas The fruit shall be produced in those areas in Xin-jiang Uygur Autonomous Region of the People's Republic of China Authorities of the People's Republic of China, as the intensive trap survey and fruit survey set forth in Article 2 below are conducted by Chinese Authorities. 2. Surveys in the production areas ① Trap survey in the designated areas during the growing season of the melon. ② Fruit survey on melon fly host plants in the designated areas during the growing season of the melon. 3. Inspection and certification in the designated production areas (a part of excerpts) In the phytosanitary certificate, the following shall bear additional remarks: The fruit has been produced within the designated production areas which are confirmed by the Chinese Authorities to be free from the Melon fly as the result of surveys set forth Article 2.	SHIP CARGO AIR CARGO															
	内モンゴル自治区 Inner Mongolia Autonomous Region 遼寧省 Liaoning 吉林省 Jilin 黒龍江省 Heilongjiang 新疆ウイグル自治区 Xinjiang Uygur Autonomous Region	PUMPKIN		May, 2008		1. Plants and areas Fresh squash shall be produced in the Neimenggu Autonomous Region, Liaoning Province, Jilin Province, Heilongjiang Province, and Xinjiang Uygur Autonomous Region of the People's Republic of China, which is designated by the Director-General of the Food Safety and Consumer Affairs Bureau of the Ministry of Agriculture, Forestry and Fisheries as an area under intensive trapping surveys and intensive fresh fruit surveys of melon flies set forth in Article 2. 2. Surveys in the production areas ① Every year, from June 1 to September 30, trapping surveys shall be carried out. ② Fresh fruit surveys. 3. Inspection and certification in the designated production areas (a part of excerpts) ① The phytosanitary certificate shall bear additional remarks that the subject fresh fruits were produced in the designated production area where no melon fly has been detected as the surveys. ② That the subject fresh fruits were packed in a place designated by the Plant Quarantine Authorities of the People's Republic of China.																
		LITCHI		April, 1994	ORIENTAL FRUIT FLY COMPLEX	After the following A treatment is done, the B treatment will be started. A Vapor Heat Treatment: The fresh Litchi shall be disinfested with saturated vapor in vapor heat treatment facilities. The temperature at the innermost fruit pulp shall be raised to 46.5℃ or higher and kept for 10 minutes. B Cold Treatment: In the cold treatment facilities, fruit pulp temperature should be lowered to 2℃ within 6 hours after vapor heat treatment and kept at the fruit pulp temperature of 2℃ for 40 hours.	SHIP CARGO AIR CARGO															
REPUBLIC OF CHILE	CHERRY	1.SWEET HEART 2.VAN 3.BING 4.LAPIN 5.LAMBERT 6.RAINIER	December, 2005	CODLING MOTH	Methyl Bromide Fumigation: ① cultivars 1 ~ 6 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>64g/m<sup>3</sup></td> <td>13.5℃ or above</td> <td>2 hours</td> <td>26.9% or less (unpacked) 19.2% or less (packed)</td> </tr> </tbody> </table> ② Or all the cultivars including cultivars 1~6 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>The production of the concentration of methyl bromide at the fumigation facilities by duration</th> </tr> </thead> <tbody> <tr> <td>64g/m<sup>3</sup></td> <td>13.5℃ or above</td> <td>2 hours or more</td> <td>95.9 or up</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	64g/m <sup>3</sup>	13.5℃ or above	2 hours	26.9% or less (unpacked) 19.2% or less (packed)	Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration	64g/m <sup>3</sup>	13.5℃ or above	2 hours or more	95.9 or up	SHIP CARGO AIR CARGO
Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																			
64g/m <sup>3</sup>	13.5℃ or above	2 hours	26.9% or less (unpacked) 19.2% or less (packed)																			
Dosage	Temperature	Duration of exposure	The production of the concentration of methyl bromide at the fumigation facilities by duration																			
64g/m <sup>3</sup>	13.5℃ or above	2 hours or more	95.9 or up																			
NEW ZEALAND	CHERRY	ALL VARIETY	December, 2005	CODLING MOTH	1. Plants and areas Cherry fresh fruits shall be produced in areas designated by New Zealand plant protection authorities to be subject to intensive trapping surveys and fresh fruit surveys for Codling moths set forth in Article 2. 2. Surveys in designated production areas ① Trapping surveys ② Fresh fruit surveys 3. Inspection and Certification in Producing Area (a part of excerpts) The phytosanitary certificate shall include the following statements: (a) The fruit shall not be infested with Codling moths. (b) The fruit shall be produced at the designated area where Codling moths have not been trapped, and be produced in the designated producing area where the number of Codling moths in the survey do not exceed 15 per trap per week on average as the result of surveys set forth in Article 2. Moreover, it shall be produced in the designated area where no Codling moths have been detected as the result of fresh fruit surveys.	SHIP CARGO AIR CARGO																
	NECTARINE	1.FANTASIA 2.RED GOLD 3.FIREBRITE	December, 1988 December, 1989		Methyl Bromide Fumigation <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>64g/m<sup>3</sup></td> <td>12℃ or above</td> <td>2 hours</td> <td>40% or less</td> </tr> </tbody> </table>	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	64g/m <sup>3</sup>	12℃ or above	2 hours	40% or less									
	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																		
64g/m <sup>3</sup>	12℃ or above	2 hours	40% or less																			
APPLE	1.GALA 2.GRANNY SMITH 3.FUJI 4.BRAEBURN 5.RED DELICIOUS 6.ROYAL GALA 7.SCIROS 8.EXCEPT ABOVE	May, 1993 July, 2007 July, 2007	①CODLING MOTH ②FIRE BLIGHT	After the following A treatment is done, the B treatment will be started. A Methyl Bromide Fumigation: ① cultivars 1~7 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>24g/m<sup>3</sup></td> <td>12℃ or above</td> <td>2 hours</td> <td>40% or less</td> </tr> </tbody> </table> ② Or all the cultivars including cultivars 1~8 <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Multiplication of Methyl Bromide concentration and fumigation time at a fumigation facility</th> </tr> </thead> <tbody> <tr> <td>24g/m<sup>3</sup></td> <td>12℃ or above</td> <td>2 hours or more</td> <td>34.2 or up</td> </tr> </tbody> </table> B Cold Treatment: At cold treatment facilities, after the fruit pulp temperature reaches 2.0℃, apple fresh fruits shall be disinfested at the fruit pulp temperature of 2.0℃ or below for 25 consecutive days.	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	24g/m <sup>3</sup>	12℃ or above	2 hours	40% or less	Dosage	Temperature	Duration of exposure	Multiplication of Methyl Bromide concentration and fumigation time at a fumigation facility	24g/m <sup>3</sup>	12℃ or above	2 hours or more	34.2 or up	SHIP CARGO AIR CARGO	
Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																			
24g/m <sup>3</sup>	12℃ or above	2 hours	40% or less																			
Dosage	Temperature	Duration of exposure	Multiplication of Methyl Bromide concentration and fumigation time at a fumigation facility																			
24g/m <sup>3</sup>	12℃ or above	2 hours or more	34.2 or up																			
REPUBLIC OF PHILIPPINES	MANGO	MANILA SUPER	July, 1975	①ORIENTAL FRUIT FLY COMPLEX ②MELON FLY	Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities by using saturated vapor at the fruit pulp temperature of 46.0℃ and more for 10 minutes.	SHIP CARGO AIR CARGO																
	PAPAYA	SOLO	April, 1994		Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities by using saturated vapor at the fruit pulp temperature of 46.0℃ and more for 70 minutes.																	
FRENCH REPUBLIC	APPLE	GOLDEN DELICIOUS	September, 1997	①MEDITERRANEAN FRUIT FLY ②CODLING MOTH ③FIRE BLIGHT	After the following A treatment is done, the B treatment will be started. A Methyl Bromide Fumigation: <table border="1"> <thead> <tr> <th>Dosage</th> <th>Temperature</th> <th>Duration of exposure</th> <th>Ratio of fruit capacity</th> </tr> </thead> <tbody> <tr> <td>30g/m<sup>3</sup></td> <td>20℃ or above</td> <td>2 hours</td> <td>49% or less</td> </tr> </tbody> </table> B Cold Treatment: At cold treatment facilities, after the fruit pulp temperature reaches 1.0℃ and keeps the same treatment at and under 1.0℃ for 50 consecutive days.	Dosage	Temperature	Duration of exposure	Ratio of fruit capacity	30g/m <sup>3</sup>	20℃ or above	2 hours	49% or less	SHIP CARGO AIR CARGO								
Dosage	Temperature	Duration of exposure	Ratio of fruit capacity																			
30g/m <sup>3</sup>	20℃ or above	2 hours	49% or less																			
FEDERATIVE REPUBLIC OF BRAZIL	MANGO	TOMMY ATKINS	September, 2004	MEDITERRANEAN FRUIT FLY	Hot Water Treatment: The fruits shall be disinfested at the hot water dip treatment facilities by using hot water which temperature is 47℃ and at the pulp temperature of 46℃ and higher for 5minutes.	SHIP CARGO AIR CARGO																
		KENT	July, 2008																			

**LIST OF BAN-LIFTED PLANT ITEMS AND KEY CONDITIONS (MEET THE STANDARDS SET BY THE MINISTER OF AGRICULTURE, FORESTRY AND FISHERIES)**

1. April, 2010

COUNTRIES / DISTRICTS	FRESH FRUITS		BAN-LIFTED YEAR	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE
	ITEMS	CULTIVARS				
SOCIALIST REPUBLIC OF VIET NAM	HYLOCEREUS UNDATUS	(Reference) DRAGON FRUIT	October, 2009	①ORIENTAL FRUIT FLY COMPLEX ②MELON FLY	Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities by using saturated vapor at the fruit pulp temperature of 46.5°C for 40 minutes after a steady increase of pulp temperature up to 43.0°C.	SHIP CARGO AIR CARGO
KINGDOM OF BELGIUM	CUCUMBER TOMATO		December, 2003	MEDITERRANEAN FRUIT FLY	1.Plants and areas Cucumbers and tomatoes shall be produced in sites (Designated Production Sites) designated by the Plant Quarantine Authority of Belgium as areas where no Mediterranean fruit fly has occurred and satisfy the following conditions: ① Cucumbers and tomatoes shall be produced at the facility (Designated Production Facilities) designated by the Plant Quarantine Authority of Belgium. ② At the monitoring areas against Mediterranean fruit fly designated by Belgium Authority, they shall be the ones produced during the period that the authority confirms with no spread of the fly after the occurrence survey mentioned in Article 2. 2.The Occurrence Survey in quarantine monitoring areas, designated production sites and designated production facilities ① Pheromone trap monitoring ② Orchard fruit monitoring 3. Inspection and Certification in Producing Area (a part of excerpts) The phytosanitary certificate shall be mentioned the following additional remarks as the result of the occurrence survey mentioned in Article 2: The fruit shall be produced at the designated production facilities where no Mediterranean fruit flies have not been trapped in the designated producing areas where no Mediterranean fruit flies have not been trapped.	SHIP CARGO AIR CARGO
REPUBLIC OF PERU	MANGO	KENT	January, 2010	MEDITERRANEAN FRUIT FLY	Hot Water Treatment: The fruits shall be disinfested at the hot water dip treatment facilities by using hot water of 47.0°C to keep the pulp temperature at 46.0°C and higher.	SHIP CARGO AIR CARGO
MALAYSIA	MANGO	HARUMANIS	May, 2008	①ORIENTAL FRUIT FLY COMPLEX ②MELON FLY	Vapor Heat Treatment: The fruits shall be disinfested at vapor heat treatment facilities by using saturated vapor at the fruit pulp temperature of 46.5°C for 20 minutes after a steady increase in the fruit pulp temperature to 43.0°C without using vapor.	SHIP CARGO AIR CARGO
REPUBLIC OF SOUTH AFRICA	SWEET ORANGE	1.WASHINGTON NAVEL 2.VALENCIA 3.TOMANGO 4.PROTEA	August, 1970	MEDITERRANEAN FRUIT FLY	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches minus 0.6°C and keeps the same temperature for 12 consecutive days.	SHIP CARGO AIR CARGO
	LEMON GRAPE FRUIT CLEMANTINE		April, 1971			
			June, 2007			
UNITED MEXICAN STATES	TOMATO		April, 2006	TABACCO BLUE MOLD	Cold Treatment: The fruits shall be disinfested at cold treatment facilities, cold treatment vessels and refrigerated shipping containers, after the fruit pulp temperature reaches 0.6°C and keeps the same temperature for 14 consecutive days.	

**Reference: TREATMENTS PROPOSED BY USA & MEXICO**

1, April, 2010

COUNTRIES / DISTRICTS	FRESH FRUITS	Directorial Notice	TARGET PESTS	TREATMENT OR OTHER MEASURES	MEANS OF CONVEYANCE																										
UNITED STATES OF AMERICA (From State of Florida)	ORANGE GRAPE FRUIT LIME (EXCLUDING PERSIAN LIME) MANGO OTHER FRESH FRUITS (EXCLUDING SOUR LEMON, CITRUS LIMON) PRODUCED IN FLORIDA WHICH ARE KNOWN TO HOST CARIBFRILIES.	June 19, 1990 (2農安第3581号)	CARIBBEAN FRUIT FLY	1. Methyl Bromide Fumigation: <table border="1"> <tr> <th>Dosage</th> <th>Duration of exposure</th> <th>Temperature</th> <th>Ratio of accommodation</th> </tr> <tr> <td>40g/m<sup>3</sup></td> <td>2 hours</td> <td>24℃ - 29℃</td> <td>20% or less</td> </tr> </table>	Dosage	Duration of exposure	Temperature	Ratio of accommodation	40g/m <sup>3</sup>	2 hours	24℃ - 29℃	20% or less	SHIP CARGO AIR CARGO																		
	Dosage	Duration of exposure	Temperature	Ratio of accommodation																											
	40g/m <sup>3</sup>	2 hours	24℃ - 29℃	20% or less																											
	ORANGE GRAPE FRUIT POMELO	June 19, 1990 (2農安第3581号) October 22, 2008 (20消安第7951号)		2. Cold Treatment (after the temperature at the innermost fruit pulp shall reach the following temperature) A) Orange, grape fruit and pomelo ① Usual treatment <table border="1"> <tr> <th>Fruit temperature</th> <th>Duration of exposure</th> </tr> <tr> <td>0.6℃(33 ℉)</td> <td>14 days</td> </tr> <tr> <td>0.8℃(33.5℉)</td> <td>16 days</td> </tr> <tr> <td>1.1℃(34 ℉)</td> <td>17 days</td> </tr> <tr> <td>1.4℃(34.5℉)</td> <td>19 days</td> </tr> <tr> <td>1.7℃(35 ℉)</td> <td>20 days</td> </tr> <tr> <td>1.9℃(35.5℉)</td> <td>22 days</td> </tr> <tr> <td>2.2℃(36 ℉)</td> <td>24 days</td> </tr> </table> ② Short term treatment <table border="1"> <tr> <th>Fruit temperature</th> <th>Duration of exposure</th> </tr> <tr> <td>0.6℃(33 ℉)</td> <td>10 days</td> </tr> <tr> <td>1.1℃(34 ℉)</td> <td>12 days</td> </tr> <tr> <td>1.7℃(35 ℉)</td> <td>14 days</td> </tr> <tr> <td>2.2℃(36 ℉)</td> <td>17 days</td> </tr> </table> Note: In case orange, grape fruit and pomelo adapt to the following conditions, the short term treatment of the above ② shall be applied to them. ① The fruits shall be produced at the areas where the occurrence density of Caribbean fruit fly is low. ② At the time of export inspection, sampling fruits shall be kept at 26.7℃ for more than 10 days. After that, fruit cutting survey shall be done. As the result of the survey, there shall be no interception of Caribbean fruit fly.	Fruit temperature	Duration of exposure	0.6℃(33 ℉)	14 days	0.8℃(33.5℉)	16 days	1.1℃(34 ℉)	17 days		1.4℃(34.5℉)	19 days	1.7℃(35 ℉)	20 days	1.9℃(35.5℉)	22 days	2.2℃(36 ℉)	24 days	Fruit temperature	Duration of exposure	0.6℃(33 ℉)	10 days	1.1℃(34 ℉)	12 days	1.7℃(35 ℉)	14 days	2.2℃(36 ℉)	17 days
	Fruit temperature	Duration of exposure																													
0.6℃(33 ℉)	14 days																														
0.8℃(33.5℉)	16 days																														
1.1℃(34 ℉)	17 days																														
1.4℃(34.5℉)	19 days																														
1.7℃(35 ℉)	20 days																														
1.9℃(35.5℉)	22 days																														
2.2℃(36 ℉)	24 days																														
Fruit temperature	Duration of exposure																														
0.6℃(33 ℉)	10 days																														
1.1℃(34 ℉)	12 days																														
1.7℃(35 ℉)	14 days																														
2.2℃(36 ℉)	17 days																														
CARAMBOLA	July 29, 1994 (6農安第4621号)		B) CARAMBOLA <table border="1"> <tr> <th>Fruit temperature</th> <th>Duration of exposure</th> </tr> <tr> <td>1.1℃(34 ℉)</td> <td>15 days</td> </tr> </table>	Fruit temperature	Duration of exposure	1.1℃(34 ℉)	15 days																								
Fruit temperature	Duration of exposure																														
1.1℃(34 ℉)	15 days																														
ORANGE GRAPE FRUIT ORO BLANCO TANGERINE POMELO	May 28, 1999 (11農産第2605号) October 22, 2008 (20消安第7951号)		3. Shipping from the quarantine administrated area ① Judging from the results of trap treatment and bait spray implemented by US plant protection services, oranges, grape fruits, oroblanco, tangelin or pomelo shall be able to be imported without disinfection at the export country or at sea, if the fruits are produced at areas (quarantine administrated areas) where no Caribbean fruit flies have been detected, and also expected to be able to keep the situation maintained. In this case, it is very essential that the production area numbers or additional declaration shall be written in Phytosanitary Certificate issued by US plant protection services. ② The above Article ① is enforced in Florida. Therefore, prior adjustment with the export side is essential in importing the subject fresh fruits.																												
MANGO	June 19, 1990 (2農安第3581号)		Hot Water Treatment: ① The fruits shall be disinfested at the hot water dip treatment facilities by using hot water at the pulp temperature of 46.1℃ or higher and the temperature must not fall below 46.0℃ for 90minutes or more. ② The size of the fruit is no more than 8 and a container includes 8 fruits and the net weight is 5kg. The average weight is 625g, but each mango must not exceed 700g.																												
UNITED MEXICAN STATE (Excluding State of Chiapas)	MANGO	February 8, 1991 (3-12) May 22, 2008 (20消安第2213号) March 31, 2009 (20消安第13407号)	FRUIT FLY (Genus : ANASTREPHA) : MEXICAN FRUIT FLY : SOUTH AMERICAN FRUIT FLY	1. Hot water dip treatment The fruits shall be disinfested at the hot water dip treatment facilities by hot water at the pulp temperature of 46.1℃ or higher, (1) Elongated and flattened types (Francis and similar shaped mangoes) shall be treated as follows: ① 375 grams or less for 65minutes ② from 375grams to 570grams for 75 minutes (2) Other varieties of mangoes shall be treated as follows: ① 500 grams or less for 75 minutes ② from 500 grams to 700 grams for 90minutes, ③ from 700 grams to 900 grams for 110 minutes 2. Forced hot air treatment The fruits shall be disinfested by hot wind of 50℃ until the pulp temperature reaches 48℃. (The gross of a mango should be 700 grams or less.)	SHIP CARGO AIR CARGO																										
	GRAPE FRUIT ORANGE	December 25, 2006 (18消安第3742号) October 7, 2008 (20消安第7289号)		1. Methyl Bromide Fumigation <table border="1"> <tr> <td>Temperature</td> <td>21.1℃ - 29.0℃ (70℉ - 85℉)</td> </tr> <tr> <td>Dosage</td> <td>40g/m<sup>3</sup></td> </tr> <tr> <td>Duration of exposure</td> <td>2 hours</td> </tr> <tr> <td>Ratio of accommodation</td> <td>less than 80%</td> </tr> <tr> <td>Export inspection</td> <td>there shall be no fruit fly of anastrepha genus.</td> </tr> </table> 2. Cold treatment (main ship or refrigerated shipping container during sailings) <table border="1"> <tr> <th>Temperature</th> <th>Duration of exposure</th> </tr> <tr> <td>0.6℃(33℉)</td> <td>18 days</td> </tr> <tr> <td>1.1℃(34℉)</td> <td>20 days</td> </tr> <tr> <td>1.7℃(35℉)</td> <td>22 days</td> </tr> <tr> <td>2.2℃(36℉)</td> <td>24 days</td> </tr> </table> Note: For a refrigerated shipping container during sailing, the consignment shall be loaded to the ship after the fruit pulp temperature of the consignment has reached the regulated treatment temperature. 3. Forced hot air treatment The fruits shall be disinfested that the fruit pulp temperature reaches 44℃ within 90 minutes or more and then keeps the same temperature and more for 100 minutes.		Temperature	21.1℃ - 29.0℃ (70℉ - 85℉)	Dosage	40g/m <sup>3</sup>	Duration of exposure	2 hours	Ratio of accommodation	less than 80%	Export inspection	there shall be no fruit fly of anastrepha genus.	Temperature	Duration of exposure	0.6℃(33℉)	18 days	1.1℃(34℉)	20 days	1.7℃(35℉)	22 days	2.2℃(36℉)	24 days						
	Temperature	21.1℃ - 29.0℃ (70℉ - 85℉)																													
Dosage	40g/m <sup>3</sup>																														
Duration of exposure	2 hours																														
Ratio of accommodation	less than 80%																														
Export inspection	there shall be no fruit fly of anastrepha genus.																														
Temperature	Duration of exposure																														
0.6℃(33℉)	18 days																														
1.1℃(34℉)	20 days																														
1.7℃(35℉)	22 days																														
2.2℃(36℉)	24 days																														
GRAPE FRUIT ( Citrus paradisi ) ORANGE ( Citrus sinensis ) MANDARIN ( Citrus reticulata ) MANGO ( Mangifera indica )	June 18, 2007 (17消安第13245号)		The shipment from designated areas: The fruits produced in the areas can be imported without any disinfection only for the subject fruits from the following areas, such as Sonora, Chihuahua, Baja California Sur, Sinaloa (restricted only 5 administrative districts, such as Ahome, El Fuerte, Choix, uasave, Sinaloa de Leyva) and an area approved as a pest free area of fruit flies of Anastrepha genus. In this case, the Work Plan (maintenance activities of pest free areas, issued phytosanitary certificate and its additional declaration, packaging seals etc.) from Mexico shall be indispensable.																												